



Dr. D. Wayne Crowe
Dr. J. O. Law*
Dr. K.J. Swanson*
Dr. T. Daws*
Dr. O. Lamykina

*denotes professional corporation

Telephone (250) 334-2445 | Fax (250) 334-2642
788 GRANT AVENUE, COURTENAY, B.C. V9N2T3

Pre-Operative Circumcision Information

1. Given Tylenol 40mg suspension 30-45 minutes prior to procedure.
2. Apply EMLA cream to base of penis one hour prior to procedure.
3. Arrive 15 minutes prior to appointment time.

Post-Operative Circumcision Information

1. The plastic bell usually comes off in 3-5 days. The string and any remaining skin will fall off at that time. Occasionally the bell will come off early. This is not a concern unless there is accompanying bleeding in which case the baby should be accessed. If the bell stays on longer than one week I would like to see the baby as well.
2. No bathing until the bell comes off. Wash around the bell cautiously and avoid pulling on it.
3. Report any excessive swelling, redness or pus like discharge.
4. Report if the baby does not void within eight hours of the procedure.
5. Use Polysporin on a small piece of gauze and place this over the tip of the penis prior to securing the diaper. Change the antibiotic gauze with every diaper change.
6. You may use Tylenol 40mg suspension every four hours if needed for irritability.
7. Text or call me if you have any concerns.

Sincerely yours,

Dr. W. Crowe, Inc
Dictated but not read

19 January 2020